

The three-body Coulomb scattering problem in discrete Hilbert-space basis representation

Z. Papp^{1,2} and S. L. Yakovlev³

¹ *Institute of Nuclear Research of the Hungarian Academy of Sciences,
P.O. Box 51, H-4001 Debrecen, Hungary*

² *Institute for Theoretical Physics, University of Graz,
Universitätsplatz 5, A-8010 Graz, Austria*

³ *Department of Mathematical and Computational Physics,
Saint-Petersburg State University,
198094 Saint-Petersburg, Petrodvoretz, Uljanovskaya Str. 1, Russia
(February 9, 2008)*

For solving the $2 \rightarrow 2, 3$ three-body Coulomb scattering problem the Faddeev-Merkuriev integral equations in discrete Hilbert-space basis representation are considered. It is shown that as far as scattering amplitudes are considered the error caused by truncating the basis can be made arbitrarily small. By this truncation also the Coulomb Green's operator is confined onto the two-body sector of the three-body configuration space and in leading order can be constructed with the help of convolution integrals of two-body Green's operators. For performing the convolution integral an integration contour is proposed that is valid for all energies, including bound-state as well as scattering energies below and above the three-body breakup threshold.

PACS number(s): 21.45.+v, 03.65.Nk, 02.30.Rz, 02.60.Nm

I. INTRODUCTION

The three-body Coulomb scattering problem is one of the most challenging long-standing problems of non-relativistic quantum mechanics. The source of the difficulties is related to the long-range character of the Coulomb potential. In standard scattering theory it is supposed that asymptotically the particles move freely what is not the case if Coulombic interactions are involved. The problem has been formulated in the Faddeev-Merkuriev theory [1] on a mathematically sound and elegant way. This theory has been developed through integral equations with connected (compact) kernels and equivalently transformed into configuration-space differential equations with asymptotic boundary conditions. However, in practical calculations, so far only the latter version of the theory has been considered, since the extremely complicated structure of the Green's operators in the kernels of Faddeev-Merkuriev integral equations has not yet allowed a direct solution.

Recently a novel method was proposed for treating the three-body Coulomb problem by solving integral equations in a discrete Hilbert-space basis representation. In particular the Faddeev-Noble equations [2] were solved in the Coulomb-Sturmian basis representation. The Coulomb Green's operators occurring in the Faddeev-Noble integral equations were connected, via Lippmann-Schwinger-type equations, to the channel-distorted Coulomb Green's operators, which can then be calculated by a convolution integral of two-body Coulomb Green's operators. The method was elaborated first for bound-state problems [3] with (repulsive) Coulomb plus nuclear potentials, then it was extended for

calculating $p-d$ scattering at energies below the breakup threshold [4]. In these calculations excellent agreements with the results of other established approaches were found and also the efficiency and accuracy of this new method were demonstrated. In a more recent work atomic bound-state problems with attractive Coulomb interactions were considered [5]; among others it showed the surprising result that in this scheme a much smaller number of angular momentum channels were needed to reach convergence than in previous methods. However, the validity of the approach along the Faddeev-Noble is restricted. Without limitations of the energy range it is strictly valid only for repulsive Coulomb interactions.

The Faddeev-Merkuriev theory is more general as it comprises all kind of dynamical situations and applies to repulsive as well as attractive Coulomb interactions. In this article we consider scattering processes which start from two-body asymptotic channels, i.e. $2 \rightarrow 2$ and $2 \rightarrow 3$ scattering. We adapt the discrete Hilbert-space basis representation to the Faddeev-Merkuriev integral equations for those processes. We provide justification for the validity of the expansion method for all energies below and above the breakup threshold and for repulsive as well as attractive long-range forces.

In the following section we recapitulate the basis elements of the integral-equation formulation of the Faddeev-Merkuriev scattering theory for long-range forces. In Sec. III we present our method for solving these equations and prove the validity of the involved expansions. We summarize the results and give some conclusion in Sec. IV.

II. FADDEEV-MERKURIEV INTEGRAL EQUATIONS

We consider three particles with masses $m_\alpha, m_\beta, m_\gamma$, where (α, β, γ) are permutations of $(1, 2, 3)$. By x_α, y_α we denote the usual Jacobi coordinates, and we will also use the notation $X = \{x_\alpha, y_\alpha\} \in \mathbf{R}^6$. The three-body kinetic-energy operator is given by the following sum of two-body kinetic energy operators

$$H^0 = h_{x_\alpha}^0 + h_{y_\alpha}^0.$$

We assume that the particles interact via (long-range) Coulomb-like potential

$$v_\alpha(x_\alpha) = \frac{e_\beta e_\gamma}{|x_\alpha|} + v_\alpha^{sr}(x_\alpha), \quad (1)$$

where e denotes the electric charge. The short-range potential $v_\alpha^{sr}(x_\alpha)$ is supposed to be square integrable, $v_\alpha^{sr}(x_\alpha) \in L_2(\mathbf{R}^3)$, and consequently it should possess the asymptotic behavior

$$v_\alpha^{sr}(x_\alpha) \sim \mathcal{O}(|x_\alpha|^{-3/2-\epsilon}), \quad \epsilon > 0,$$

as $|x_\alpha| \rightarrow \infty$.

The Faddeev integral equations are the fundamental equations for three-body problems. They were originally formulated for short-range interactions [6]. In this case, after one iteration they possess connected kernels, and consequently they are effectively Fredholm-type integral equations of the second kind. The fact that the homogeneous equations allow nontrivial solutions only for a discrete set of three-body bound-state energies guarantees the existence of unique solutions of the inhomogeneous equations at scattering energies. For long-range interactions, however, the integral kernels of the Faddeev equations become singular and the nice properties above do no longer hold. Therefore one cannot simply plug in a Coulomb-like potential into the Faddeev equations.

A natural way of remedy is to split somehow the Coulomb-like interaction into short-range and long-range terms

$$v_\alpha = v_\alpha^{(s)} + v_\alpha^{(l)},$$

and apply the Faddeev procedure only to the short-range parts. Here, the superscripts s and l indicate the short- and long-range attributes, respectively. The three-body Hamiltonian is defined by

$$H = H^0 + v_\alpha^{(s)} + v_\alpha^{(l)} + v_\beta^{(s)} + v_\beta^{(l)} + v_\gamma^{(s)} + v_\gamma^{(l)}.$$

Merkuriev introduced the splitting in the three-body configuration space $\mathbf{R}^6$ via a splitting function ζ_α ,

$$v_\alpha^{(s)}(X) = v_\alpha(x_\alpha)\zeta_\alpha(x_\alpha, y_\alpha),$$

$$v_\alpha^{(l)}(X) = v_\alpha(x_\alpha)[1 - \zeta_\alpha(x_\alpha, y_\alpha)].$$

The function ζ_α is defined such that it separates the asymptotic two-body sector Ω_α from the rest of the three-body configuration space. On the region of Ω_α the splitting function ζ_α asymptotically tends to 1 and on the complementary asymptotic region of the configuration space it tends to 0. Rigorously, Ω_α is defined as a part of the three-body configuration space where the condition

$$|x_\alpha| < a(1 + |y_\alpha|/a)^\nu, \text{ with } a > 0, 0 < \nu < 1/2, \quad (2)$$

is satisfied. So, in Ω_α the short-range part $v_\alpha^{(s)}$ coincides with the original Coulomb-like potential v_α and in the complementary region vanishes, whereas the opposite holds true for $v_\alpha^{(l)}$.

We define the long-ranged and the channel long-ranged Hamiltonians as

$$H^{(l)} = H^0 + v_\alpha^{(l)} + v_\beta^{(l)} + v_\gamma^{(l)},$$

and

$$H_\alpha^{(l)} = H^0 + v_\alpha^{(s)} + v_\alpha^{(l)} + v_\beta^{(l)} + v_\gamma^{(l)},$$

respectively, together with their resolvent operators

$$G^{(l)}(z) = (z - H^{(l)})^{-1}$$

and

$$G_\alpha^{(l)}(z) = (z - H_\alpha^{(l)})^{-1}.$$

The Hamiltonian $H^{(l)}$ is constructed to be a positive operator with continuous spectrum $\sigma(H^{(l)}) = [0, \infty)$ [7,8]. This continuous spectrum is structureless, its properties are similar to the spectrum of the three-body kinetic energy operator. As far as the resolvent is concerned it means that $G^{(l)}(z)$ is an analytic operator-valued function in the complex z plane except for the cut coinciding with $\sigma(H^{(l)})$. This is a very important property, it ensures that the Faddeev procedure becomes an asymptotic filtering [9] and consequently guarantees the asymptotic orthogonality of the Faddeev components. In Noble's approach [2] the splitting is performed in the two-body configuration space, like in Eq. (1), thus the above condition, without any restriction to the energy range, can be fulfilled only in case of repulsive Coulomb interactions. Now, if we perform the Faddeev decomposition of the three-body wave function $|\Psi\rangle$,

$$|\psi_\alpha\rangle = G^{(l)}v_\alpha^{(s)}|\Psi\rangle,$$

it will play the role of an asymptotic filtering. It leads to the decomposition of the wave function into three components

$$|\Psi\rangle = |\psi_\alpha\rangle + |\psi_\beta\rangle + |\psi_\gamma\rangle,$$

with each of them describing different asymptotic channels. The resulting equations for the components are the set of Faddeev-Merkuriev integral equations [8]

$$|\psi_\alpha\rangle = \delta_{\beta\alpha}|\Phi_{\beta i}^{(l)+}\rangle + G_\alpha^{(l)}(E + i0)v_\alpha^{(s)} \sum_{\gamma \neq \alpha} |\psi_\gamma\rangle, \quad (3)$$

where $|\Phi_{\beta i}^{(l)+}\rangle$ is an eigenstate of $H_\beta^{(l)}$ with energy E . Since we consider here only $2 \rightarrow 2, 3$ processes $|\Phi_{\beta i}^{(l)+}\rangle$ describes asymptotically bound-state pairs in the two-body subsystem β with binding energy $E_{\beta i}$. Merkuriev has shown in Ref. [7,8] that after a certain number of iterations Eqs. (3) are reduced to Fredholm integral equations of the second kind with compact kernels for all energies, including energies below ($E < 0$) and above ($E > 0$) the three-body breakup threshold. Thus all the nice properties of the original Faddeev equations established for short-range interactions remain valid also for the case of Coulomb-like potentials.

It is important to note here that the resolvent $G_\alpha^{(l)}$ of the channel long-ranged Hamiltonian $H_\alpha^{(l)}$ is an operator which, in contrast to the short-range case, cannot be constructed only from two-body quantities. To determine $G_\alpha^{(l)}$ uniquely one should start again from Faddeev-type resolvent equations, or from the triad of Lippmann-Schwinger equations [10], which is, in fact, the adjoint representation of the Faddeev operator [11]. The Hamiltonian $H_\alpha^{(l)}$, however, has the advantageous property that it supports bound states only in the two-body subsystem α , and thus only one kind of asymptotic channel is possible, the α channel. For such systems one single Lippmann-Schwinger equation treating properly the long-range part of $H_\alpha^{(l)}$ is sufficient for a unique solution [12]. The appropriate Lippmann-Schwinger equation was proposed by Merkuriev and has the form

$$G_\alpha^{(l)}(z) = G_\alpha^{as}(z) + G_\alpha^{as}(z)V_\alpha^{as}G_\alpha^{(l)}(z), \quad (4)$$

with a three-body potential V_α^{as} , which, as $|X| \rightarrow \infty$, decays faster than the Coulomb potential in all direction of the three-body configuration space $\mathbf{R}^6$: $V_\alpha^{as} \sim \mathcal{O}(|X|^{-1-\epsilon})$, $\epsilon > 0$.

The operators G_α^{as} and V_α^{as} are complicated three-body operators. However, in the region Ω_α they take relatively simple forms. The potential V_α^{as} is defined by

$$V_\alpha^{as} \equiv U^\alpha = v_\beta^{(l)} + v_\gamma^{(l)} - u_\alpha^{(l)}, \quad (5)$$

where the auxiliary two-body potential $u_\alpha^{(l)}(y_\alpha)$ must be taken in such a way that it has the asymptotic form $u_\alpha^{(l)}(y_\alpha) \sim e_\alpha(e_\beta + e_\gamma)/|y_\alpha|$, as $|y_\alpha| \rightarrow \infty$. In this case U^α , as $|y_\alpha| \rightarrow \infty$, decays sufficiently fast, namely

$$U^\alpha \sim \mathcal{O}(|y_\alpha|^{-2+\nu}), \quad (6)$$

which, considering (2), means square integrability. So, in the two-body region Ω_α the auxiliary long-range potential $u_\alpha^{(l)}$ compensates asymptotically the long-range tail of $v_\beta^{(l)} + v_\gamma^{(l)}$.

In this region also the Green's operator $G_\alpha^{as}(z)$ takes a simple form, it coincides with the channel distorted

Coulomb Green's operator $\tilde{G}_\alpha(z)$. The operator $\tilde{G}_\alpha(z)$ appears as a resolvent of the sum of two commuting two-body Hamiltonians

$$\tilde{G}_\alpha(z) = (z - h_{x_\alpha} - h_{y_\alpha})^{-1},$$

where $h_{x_\alpha} = h_{x_\alpha}^0 + v_\alpha$ and $h_{y_\alpha} = h_{y_\alpha}^0 + u_\alpha^{(l)}$, therefore it can be constructed as a convolution integral of two-body Green's operators

$$G_\alpha^{as}(z) \equiv \tilde{G}_\alpha(z) = \frac{1}{2\pi i} \oint_C dz' g_{x_\alpha}(z - z') g_{y_\alpha}(z'), \quad (7)$$

where $g_{x_\alpha}(z) = (z - h_{x_\alpha})^{-1}$ and $g_{y_\alpha}(z) = (z - h_{y_\alpha})^{-1}$. The contour C is to be taken counterclockwise around the continuous spectrum of h_{y_α} so that g_{x_α} is analytic on the domain encircled by C .

The spectra of $G_\alpha^{(l)}$ and of G_α^{as} should be of similar nature. Therefore, if we want that $\tilde{G}_\alpha$ replaces G_α^{as} on the domain of Ω_α it must possess an analogous spectrum. Since $G_\alpha^{(l)}$ has been constructed such that the branch points of its spectrum are related to the two-body bound states generated by $v_\alpha(x_\alpha)$ the auxiliary two-body potential $u_\alpha^{(l)}(y_\alpha)$ must be taken in such a way that it does not support any bound state. If $e_\alpha(e_\beta + e_\gamma) \geq 0$, it is easy to fulfill this requirement, if $e_\alpha(e_\beta + e_\gamma) < 0$ the infinitely many bound states should be projected out leading to a nonlocal $u_\alpha^{(l)}$.

All these properties of Eq. (4) lead to the following asymptotic behavior of the integral

$$F_\alpha^{(l)}(X) = \int dX' G_\alpha^{(l)}(X, X', E + i0) f(X') \sim Q_0(X, E) \langle \Phi_{\alpha 0}^{(l)-} | f \rangle + \sum_j \phi_{\alpha j}(x_\alpha) Q_{\alpha j}(y_\alpha, E) \langle \Phi_{\alpha j}^{(l)-} | f \rangle, \quad (8)$$

where $f \in L_2$. Here $\phi_{\alpha j}(x_\alpha)$ is the j -th bound-state wave function of the pair α with energy $E_{\alpha j}$. The state $|\Phi_{\alpha 0}^{(l)}\rangle$ is a scattering eigenstate of the Hamiltonian $H_\alpha^{(l)}$ describing in the final state a Coulomb distorted propagation of the three particles. The state $|\Phi_{\alpha j}^{(l)}\rangle$ describes the scattering of the third particle with respect to the bound pair. Its asymptotic form is given as a product of two-body wave functions

$$\langle x_\alpha y_\alpha | \Phi_{\alpha j}^{(l)} \rangle \sim \langle x_\alpha | \phi_{\alpha j} \rangle \langle y_\alpha | \chi_\alpha^{(l)} \rangle,$$

where $|\chi_\alpha^{(l)}\rangle$ is a scattering eigenstate of h_{y_α} . The Coulomb spherical wave functions possess the asymptotic behavior $Q_0(X, E) \sim |X|^{-5/2}$, as $|X| \rightarrow \infty$, and $Q_{\alpha j}(y_\alpha, E) \sim |y_\alpha|^{-1}$, as $|y_\alpha| \rightarrow \infty$, respectively [1]. From its construction it follows that the state $|F_\alpha^{as}\rangle = G_\alpha^{as}|f\rangle$ possesses an asymptotic behavior similar to $|F_\alpha^{(l)}\rangle$ of Eq. (8).

The asymptotics (8) leads immediately to an analogous asymptotic form for the solutions to the Faddeev-Merkuriev equations

$$\begin{aligned} \psi_\alpha(X) \sim & \delta_{\beta\alpha} \phi_{\beta i}(x_\beta) \chi_\beta^{(l)}(y_\beta, E - E_{\beta i}) + \\ & \sum_j \phi_{\alpha j}(x_\alpha) Q_{\alpha j}(y_\alpha, E) A_{\alpha j, \beta i} + \\ & Q_0(X, E) A_{\alpha 0, \beta i}, \end{aligned} \quad (9)$$

where the scattering amplitudes are defined by

$$A_{\alpha j, \beta i} = \sum_{\gamma \neq \alpha} \langle \Phi_{\alpha j}^{(l)-} | v_\alpha^{(s)} | \psi_\gamma \rangle \quad (10)$$

and

$$A_{\alpha 0, \beta i} = \sum_{\gamma \neq \alpha} \langle \Phi_{\alpha 0}^{(l)-} | v_\alpha^{(s)} | \psi_\gamma \rangle. \quad (11)$$

Finally we should note that the term $v_\alpha^{(s)} | \psi_\gamma \rangle$, $\gamma \neq \alpha$, appearing at the right hand side of Eq. (3) is a square integrable function in the three-body Hilbert space: $v_\alpha^{(s)} | \psi_\gamma \rangle \in L_2(\mathbf{R}^6)$. In fact, according to (9), we need to consider only terms like $v_\alpha^{(s)} \phi_{\gamma j}(x_\gamma)$ and $v_\alpha^{(s)} Q_0(X, E)$. The function $v_\alpha^{(s)} \phi_{\gamma j}(x_\gamma)$ is square integrable due to the fast decrease of the potential $v_\alpha^{(s)}$ in the coordinate x_α and of the two-body bound-state wave function $\phi_{\gamma j}(x_\gamma)$ in the coordinate x_γ . In the term $v_\alpha^{(s)} Q_0(X, E)$ the potential $v_\alpha^{(s)}$ contains the splitting function ζ_α , which restricts the domain of integration to Ω_α . On this domain, however, $Q_0(X, E)$ itself is square integrable if $\nu < 1/2$, therefore $v_\alpha^{(s)} Q_0(X, E)$ is also square integrable in $\mathbf{R}^6$.

III. SOLUTION OF THE INTEGRAL EQUATIONS

In this section we consider a discrete Hilbert space basis representation of the Faddeev-Merkuriev integral equations. We will show that the truncation of the basis at finite N leads to a negligible error in the scattering amplitudes if N tends to infinity.

A. Discrete Hilbert space basis representation of the Faddeev-Merkuriev integral equations

We suppose that the states $\{|n\rangle\}$ together with their biorthogonal partner $\{|\tilde{n}\rangle\}$ form a basis in the Hilbert space $L_2(\mathbf{R}^3)$ of the two-body relative motion. Since the three-body Hilbert space is a tensor product of two such two-body spaces, an appropriate basis is the direct product

$$|nm\rangle_\alpha = |n\rangle_\alpha \otimes |m\rangle_\alpha, \quad (n, m = 0, 1, 2, \dots). \quad (12)$$

The completeness of the basis in the three-body Hilbert space $L_2(\mathbf{R}^6)$ can be expressed in the form

$$\mathbf{1} = \lim_{N \rightarrow \infty} \sum_{n, m=0}^N |\widetilde{nm}\rangle_\alpha \langle nm| = \lim_{N \rightarrow \infty} \mathbf{1}_\alpha^N.$$

Note that, in the three-body Hilbert space, three equivalent bases belonging to fragmentations α , β , and γ are possible.

We start from the observation of the previous section that each term $v_\alpha^{(s)} | \psi_\gamma \rangle$, $\alpha \neq \gamma$, in Eqs. (3) is a square integrable function and consequently can be expanded in $L_2(\mathbf{R}^6)$ in the convergent series

$$\begin{aligned} v_\alpha^{(s)} | \psi_\gamma \rangle &= \sum_{n, m=0}^{\infty} |\widetilde{nm}\rangle_\alpha \langle nm | v_\alpha^{(s)} | \psi_\gamma \rangle \\ &= \mathbf{1}_\alpha^{N_1} v_\alpha^{(s)} | \psi_\gamma \rangle + |r_\alpha^{N_1}\rangle. \end{aligned} \quad (13)$$

In the last line of this equation we have truncated the series at some finite N_1 and represented the remainder as $|r_\alpha^{N_1}\rangle = (\mathbf{1} - \mathbf{1}_\alpha^{N_1}) v_\alpha^{(s)} | \psi_\gamma \rangle$. Due to the square integrability of $v_\alpha^{(s)} | \psi_\gamma \rangle$ the latter has the zero limit

$$\lim_{N_1 \rightarrow \infty} \|r_\alpha^{N_1}\| = 0, \quad (14)$$

where $\|f\| = \langle f | f \rangle^{1/2}$ stands for the Hilbert-space norm. Now, the Faddeev-Merkuriev integral equations (3) take the form

$$\begin{aligned} |\psi_\alpha\rangle &= \delta_{\alpha\beta} |\Phi_{\beta i}^{(l)+}\rangle + G_\alpha^{(l)}(E + i0) \mathbf{1}_\alpha^{N_1} v_\alpha^{(s)} \sum_{\gamma \neq \alpha} |\psi_\gamma\rangle \\ &\quad + G_\alpha^{(l)}(E + i0) |r_\alpha^{N_1}\rangle. \end{aligned} \quad (15)$$

To proceed further let us consider the terms $\langle nm | v_\alpha^{(s)} | \psi_\gamma \rangle$ appearing in the second term on r.h.s. of (15). This matrix element can explicitly be expressed by the integral

$$\langle nm | v_\alpha^{(s)} | \psi_\gamma \rangle = \int dx_\alpha dy_\alpha \langle nm | X \rangle v_\alpha^{(s)}(X) \psi_\gamma(X). \quad (16)$$

The product $\langle nm | X \rangle v_\alpha^{(s)}(X)$ in the integral is square integrable and again can be expanded in $L_2(\mathbf{R}^6)$ using the basis (12) in fragmentation γ

$$\begin{aligned} \langle nm | X \rangle v_\alpha^{(s)}(X) &= \sum_{n', m'}^{\infty} \langle nm | v_\alpha^{(s)} | n' m' \rangle_\gamma \langle \widetilde{n' m'} | X \rangle \\ &= \langle nm | v_\alpha^{(s)} \mathbf{1}_\gamma^{N_2} | X \rangle + \\ &\quad \langle nm | v_\alpha^{(s)} (\mathbf{1} - \mathbf{1}_\gamma^{N_2}) | X \rangle. \end{aligned} \quad (17)$$

Again the expansion is terminated at some finite N_2 , but this is chosen such that for all $n, m \leq N_1$ the remainder becomes arbitrarily small:

$$\| \langle nm | v_\alpha^{(s)} (\mathbf{1} - \mathbf{1}_\gamma^{N_2}) \| < \varepsilon; \quad \varepsilon > 0. \quad (18)$$

Substituting Eq. (17) into Eq. (15) we arrive at the following representation

$$|\psi_\alpha\rangle = \delta_{\alpha\beta}|\Phi_{\beta i}^{(l)+}\rangle + G_\alpha^{(l)}(E+i0)\mathbf{1}_\alpha^{N_1}v_\alpha^{(s)}\sum_{\gamma\neq\alpha}\mathbf{1}_\gamma^{N_2}|\psi_\gamma\rangle \\ + G_\alpha^{(l)}(E+i0)[|r_\alpha^{N_1}\rangle + |r_\alpha^{N_1N_2}\rangle], \quad (19)$$

where $|r_\alpha^{N_1N_2}\rangle = \mathbf{1}_\alpha^{N_1}v_\alpha^{(s)}\sum_{\gamma\neq\alpha}(\mathbf{1}-\mathbf{1}_\gamma^{N_2})|\psi_\gamma\rangle$.

Below we will show that the last term in Eqs. (19) gives a negligible contribution to the scattering amplitudes if N_1 and N_2 tend to infinity. Indeed, from Eqs. (14) and (18) it follows that for any $\varepsilon > 0$ there exist N_1 and N_2 such that for all $N \geq \max\{N_1, N_2\}$ the inequality

$$\|r_\alpha^N + r_\alpha^{NN}\| < \varepsilon$$

holds. The last term on the right-hand side of Eqs. (19) can now be considered as an additional driving term. Then the solution of Eqs. (19) can be represented as a sum of two terms

$$|\psi_\alpha\rangle = |\psi_\alpha^N\rangle + |R_\alpha^N\rangle, \quad (20)$$

which satisfy the Faddeev-Merkuriev integral equations with degenerate kernels of rank N

$$|\psi_\alpha^N\rangle = \delta_{\alpha\beta}|\Phi_{\beta i}^{(l)+}\rangle + G_\alpha^{(l)}(E+i0)\mathbf{1}_\alpha^N v_\alpha^{(s)}\sum_{\gamma\neq\alpha}\mathbf{1}_\gamma^N|\psi_\gamma^N\rangle \quad (21)$$

$$|R_\alpha^N\rangle = G_\alpha^{(l)}(E+i0)[|r_\alpha^N\rangle + |r_\alpha^{NN}\rangle] + \\ G_\alpha^{(l)}(E+i0)\mathbf{1}_\alpha^N v_\alpha^{(s)}\sum_{\gamma\neq\alpha}\mathbf{1}_\gamma^N|R_\gamma^N\rangle. \quad (22)$$

Due to the degeneracy of the kernels these equations are automatically compact. Actually, only the first one is needed for practical calculations because the solution of the second one can be made arbitrarily small by a proper choice of the parameter N . To show this explicitly we rewrite Eqs. (22) in differential form

$$(E - H_\alpha^{(l)})|R_\alpha^N\rangle - \mathbf{1}_\alpha^N v_\alpha^{(s)}\sum_{\gamma\neq\alpha}\mathbf{1}_\gamma^N|R_\gamma^N\rangle = |r_\alpha^N\rangle + |r_\alpha^{NN}\rangle,$$

and express its solution via the resolvent of the Faddeev matrix operator [11]

$$|R_\alpha^N\rangle = \sum_\gamma G_{\alpha\gamma}^N(E+i0)(|r_\gamma^N\rangle + |r_\gamma^{NN}\rangle).$$

The resolvent components $G_{\alpha\gamma}^N$ are defined as the solution of the Faddeev-Merkuriev integral equations

$$G_{\beta\alpha}^N(z) = G_\beta^{(l)}(z)\delta_{\beta\alpha} + G_\beta^{(l)}(z)\mathbf{1}_\beta^N v_\beta^{(s)}\sum_{\gamma\neq\beta}\mathbf{1}_\gamma^N G_{\gamma\alpha}^N(z),$$

which, as before, are automatically compact due to the degeneracy of the kernel. The same arguments that allowed to arrive at the asymptotics (8) lead to a similar asymptotic form for the kernel $G_{\beta\alpha}^N(X, X', E+i0)$. One needs only to replace the states $|\Phi_{\alpha j}^{(l)-}\rangle$ and $|\Phi_{\alpha 0}^{(l)-}\rangle$ by the

three-body wave functions $|\Psi_{\alpha j}^-\rangle$ and $|\Psi_{\alpha 0}^-\rangle$, respectively. This involves a similar asymptotic form of R_α^N as for ψ_α in Eq. (9) with amplitudes

$$B_{\alpha j}^N = \sum_\gamma \langle \Psi_{\alpha j}^- | (|r_\gamma^N\rangle + |r_\gamma^{NN}\rangle) \rangle$$

and

$$B_{\alpha 0}^N = \sum_\gamma \langle \Psi_{\alpha 0}^- | (|r_\gamma^N\rangle + |r_\gamma^{NN}\rangle) \rangle.$$

From the unitarity relation of the three-body wave functions we get the inequality

$$\sum_{\alpha j} |B_{\alpha j}^N|^2 + \sum_\alpha |B_{\alpha 0}^N|^2 = \sum_\gamma \|r_\gamma^N + r_\gamma^{NN}\|^2 < 3\varepsilon^2.$$

This means that with a proper choice of N the moduli of the amplitudes can be made arbitrarily small and the neglect of $|R_\alpha^N\rangle$ in (20) causes only a negligible error in the scattering amplitudes.

So far we have reduced the problem of solving Eqs. (3) to the solution of Eqs. (21) with a degenerate kernel for $|\psi_\alpha^N\rangle$. Its asymptotic behavior is exactly the same as of the original equations (9), only the amplitudes $A_{\alpha j, \beta i}$ and $A_{\alpha j, \beta i}$ have to be replaced by

$$A_{\alpha j, \beta i}^N = \sum_{\gamma\neq\alpha} \langle \Phi_{\alpha j}^{(l)-} | v_\alpha^{(s)} | \psi_\gamma^N \rangle$$

and

$$A_{\alpha 0, \beta i}^N = \sum_{\gamma\neq\alpha} \langle \Phi_{\alpha 0}^{(l)-} | v_\alpha^{(s)} | \psi_\gamma^N \rangle,$$

respectively. So, in terms of amplitudes A and A^N we can state that for any $\varepsilon > 0$ there exists N_0 such that for every $N > N_0$ the relation

$$\sum_{\alpha j} |A_{\alpha j, \beta i} - A_{\alpha j, \beta i}^N|^2 + \sum_\alpha |A_{\alpha 0, \beta i} - A_{\alpha 0, \beta i}^N|^2 < 3\varepsilon^2$$

holds, i.e. $A_{\alpha j, \beta i}^N$ and $A_{\alpha 0, \beta i}^N$ as N goes to infinity tend to $A_{\alpha j, \beta i}$ and $A_{\alpha 0, \beta i}$, respectively.

In calculating $A_{\alpha j, \beta i}^N$ and $A_{\alpha 0, \beta i}^N$ we can again perform an approximation. Recalling the fact that $v_\alpha^{(s)}|\psi_\gamma^N\rangle$ is square integrable, similarly to the approximations made before in formulas (13) and (17), we perform approximations to the scattering amplitudes

$$A_{\alpha j, \beta i}^{NM} = \sum_{\gamma\neq\alpha} \langle \Phi_{\alpha j}^{(l)-} | \mathbf{1}_\alpha^M v_\alpha^{(s)} \mathbf{1}_\gamma^M | \psi_\gamma^N \rangle \quad (23)$$

and

$$A_{\alpha 0, \beta i}^{NM} = \sum_{\gamma\neq\alpha} \langle \Phi_{\alpha 0}^{(l)-} | \mathbf{1}_\alpha^M v_\alpha^{(s)} \mathbf{1}_\gamma^M | \psi_\gamma^N \rangle, \quad (24)$$

which, as M goes to infinity tend to $A_{\alpha j, \beta i}^N$ and $A_{\alpha 0, \beta i}^N$, respectively. Summarizing our results we can conclude that

$$\sum_{\alpha j} |A_{\alpha j, \beta i} - A_{\alpha j, \beta i}^{NM}|^2 + \sum_{\alpha} |A_{\alpha 0, \beta i} - A_{\alpha 0, \beta i}^{NM}|^2 \rightarrow 0,$$

as N and M go to infinity. Thus the reduction of the Faddeev-Merkuriev integral equations to matrix equations and the use of the resulting finite basis representation of the Faddeev components in calculating scattering amplitudes via (23) and (24) is justified, the error made by using finite N and M values in the truncation can be made arbitrarily small.

In practical calculations all what we need is to solve the matrix equations for $\underline{\psi}_\alpha = {}_\alpha \langle \widetilde{nm} | \psi_\alpha \rangle$

$$\underline{\psi}_\alpha = \delta_{\alpha\beta} \underline{\Phi}_{\beta i}^{(l)+} + \underline{G}_\alpha^{(l)} \sum_{\gamma \neq \alpha} \underline{v}_{\alpha\gamma}^{(s)} \underline{\psi}_\gamma,$$

where $\underline{G}_\alpha^{(l)} = {}_\alpha \langle \widetilde{nm} | G_\alpha^{(l)} | \widetilde{n'm'} \rangle_\alpha$, $\underline{v}_{\alpha\gamma}^{(s)} = {}_\alpha \langle nm | v_\alpha^{(s)} | n'm' \rangle_\gamma$ and $\underline{\Phi}_\alpha^{(l)} = {}_\alpha \langle \widetilde{nm} | \Phi_\alpha^{(l)} \rangle$, and calculate the scattering amplitudes by

$$A_{\alpha j, \beta i} = \sum_{\gamma \neq \alpha} \underline{\Phi}_{\alpha j}^{(l)-} \underline{v}_{\alpha\gamma}^{(s)} \underline{\psi}_\gamma$$

and

$$A_{\alpha 0, \beta i} = \sum_{\gamma \neq \alpha} \underline{\Phi}_{\alpha 0}^{(l)-} \underline{v}_{\alpha\gamma}^{(s)} \underline{\psi}_\gamma.$$

B. Evaluation of matrix elements $\underline{G}_\alpha^{(l)}$ and $\underline{\Phi}_\alpha^{(l)}$

While the matrix elements of the potential $v_\alpha^{(s)}$ can always be calculated, at least numerically, we need further approximations to calculate the L_2 matrix elements $\underline{G}_\alpha^{(l)}$ and $\underline{\Phi}_\alpha^{(l)}$.

For calculating $\underline{G}_\alpha^{(l)}$ we make use of the Lippmann-Schwinger equation (4), and we show that this integral equation for the matrix elements can be reduced to an integral equation with a degenerate kernel. Eq. (4) for $\underline{G}_\alpha^{(l)}$ takes the form

$$\underline{G}_\alpha^{(l)} = \langle \widetilde{nm} | G_\alpha^{as} | \widetilde{n'm'} \rangle + \langle \widetilde{nm} | G_\alpha^{as} V_\alpha^{as} G_\alpha^{(l)} | \widetilde{n'm'} \rangle. \quad (25)$$

Let us consider a typical integral appearing in the second term of the previous equation

$$\langle f | G_\alpha^{as} V_\alpha^{as} G_\alpha^{(l)} | f' \rangle = \int dy''_\alpha dx''_\alpha F_\alpha^{as}(X'') V_\alpha^{as}(X'') F_\alpha^{(l)}(X''), \quad (26)$$

where $f, f' \in L_2$, and $F_\alpha^{(l)}$ and F_α^{as} are defined in (8). We split the integration domain into two parts as

$$\int_{\mathbf{R}^6} dX'' = \int_{|x''_\alpha| \leq \xi} dX'' + \int_{|x''_\alpha| > \xi} dX''.$$

The leading term of the second integral, due to (8), for large ξ values takes the form

$$I(\xi) \sim \int_0^\infty dy y^2 \int_\xi^\infty dx x^2 \frac{\exp\{i2\sqrt{E}\sqrt{x^2+y^2}\}}{(x^2+y^2)^{(6+\epsilon)/2}},$$

and the asymptotic analysis, which involves integration by parts, results

$$I(\xi) \sim \mathcal{O}(\xi^{-1-\epsilon}).$$

In other words, the integral (26) can be confined to the asymptotic two-body sector Ω_α of the configuration space. Thus we can use the explicit forms of G_α^{as} and V_α^{as} given by (7) and (5) valid on this domain and instead of (25) we can use

$$\underline{G}_\alpha^{(l)} = \langle \widetilde{nm} | \widetilde{G}_\alpha | \widetilde{n'm'} \rangle + \langle \widetilde{nm} | \widetilde{G}_\alpha U^\alpha G_\alpha^{(l)} | \widetilde{n'm'} \rangle,$$

where the restriction of integration on domain Ω_α is assumed implicitly. Now, due to the square-integrability of U^α with respect to the variable y_α (see Eq. (6)) and because of Eq. (8), we can write

$$U^\alpha G_\alpha^{(l)} | \widetilde{n'm'} \rangle = \mathbf{1}_\alpha^N U^\alpha G_\alpha^{(l)} | \widetilde{n'm'} \rangle + (\mathbf{1} - \mathbf{1}_\alpha^N) U^\alpha G_\alpha^{(l)} | \widetilde{n'm'} \rangle.$$

Repeating the arguments that lead us to Eqs. (19) we get

$$\begin{aligned} \underline{G}_\alpha^{(l)} &= \widetilde{G}_\alpha + \widetilde{G}_\alpha U^\alpha \underline{G}_\alpha^{(l)} + \\ &\quad \langle \widetilde{nm} | \widetilde{G}_\alpha (\mathbf{1} - \mathbf{1}_\alpha^N) U^\alpha G_\alpha^{(l)} | \widetilde{n'm'} \rangle + \\ &\quad \langle \widetilde{nm} | \widetilde{G}_\alpha \mathbf{1}_\alpha^N U^\alpha (\mathbf{1} - \mathbf{1}_\alpha^N) G_\alpha^{(l)} | \widetilde{n'm'} \rangle. \end{aligned}$$

By the same argumentation as before one can easily show that the contribution of the last two terms to the solution $\underline{G}_\alpha^{(l)}$ can be made arbitrarily small by a proper choice of the expansion parameter N . Thus, we have reduced the problem of solving Eq. (25) to the purely algebraic problem

$$\underline{G}_\alpha^{(l)} = \widetilde{G}_\alpha + \widetilde{G}_\alpha U^\alpha \underline{G}_\alpha^{(l)}.$$

A similar equation holds true for the wave function matrix elements

$$\underline{\Phi}_\alpha^{(l)} = \widetilde{\Phi}_\alpha + \widetilde{G}_\alpha U^\alpha \underline{\Phi}_\alpha^{(l)},$$

where $\widetilde{\Phi}_\alpha = \langle \widetilde{n} | \phi_\alpha \rangle \otimes \langle \widetilde{m} | \chi_\alpha^{(l)} \rangle$ are direct products of two-body wave function matrix elements.

For evaluating the matrix elements $\widetilde{G}_\alpha$ we can use the contour integral representation (7)

$$\widetilde{G}_\alpha(z) = \frac{1}{2\pi i} \oint_C dz' {}_\alpha \langle \widetilde{n} | g_{x_\alpha}(z-z') | \widetilde{n'} \rangle_\alpha {}_\alpha \langle \widetilde{m} | g_{y_\alpha}(z') | \widetilde{m'} \rangle_\alpha.$$

The contour C is to be chosen so that it encircles the spectrum of g_{y_α} and avoids the singularities of g_{x_α} . For bound-state energies the spectra of the two Green's operators are well separated and this condition can be fulfilled easily [3]. Also, for scattering energies below breakup

threshold, where the bound-state pole of g_{x_α} can meet the continuous spectrum of g_{y_α} , the contour integration can still be performed [4]. For other positive-energy scattering problems, however, the continua overlap and therefore these contours are not viable.

Fortunately, there exists a contour that is valid for all z of physical interest. Besides positive real values of z that arise for scattering above the breakup threshold, this contour can deal with complex values of z having negative imaginary parts that are needed for resonant-state calculations. In this approach one must first shift the spectrum of g_{x_α} by taking $z = E + i\varepsilon$ with positive ε . By doing so, the two spectra become well separated and the spectrum of g_{y_α} can be encircled. Next the contour C is deformed analytically in such a way that the upper part descends to the second Riemann sheet of g_{y_α} , while the lower part of C can be turned away from the cut [see Fig. 1]. The contour still encircles the branch cut singularity of g_{y_α} , but in the limit $\varepsilon \rightarrow 0$ it now avoids the singularities of g_{x_α} . Moreover, by continuing to negative values of ε , the branch cut and pole singularities of g_{x_α} move to the second Riemann sheet of g_{y_α} and, at the same time, the branch cut of g_{y_α} moves to the second Riemann sheet of g_{x_α} . Thus, the mathematical conditions for the contour integral representation of $\tilde{G}_\alpha(z)$ in Eq. (7) can be fulfilled for all energies. In this respect there is only a gradual difference between the calculations for bound and resonant states as well as scattering below and above the breakup threshold.

IV. SUMMARY

In this paper we have shown that in the kernels of the Faddeev-Merkuriev integral equations for $2 \rightarrow 2, 3$ scattering the short-range potentials can be approximated with the use of a discrete Hilbert-space basis. Then, to solve for scattering amplitudes one needs only the matrix elements of the channel long-ranged Green's operator between basis states. For evaluating these matrix elements we use specific Lippmann-Schwinger equation, and we show that in this particular case the potential term in the kernel can again be approximated on the same finite basis. This truncation involves a reduction of the asymptotic channel long-ranged Green's operator onto the two-body sector of the three-body configuration space, where it can be represented in leading order by a convolution integral of two-body Green's operators. For performing the convolution integral we propose a contour that applies for all energies of physical interest.

In principle any discrete Hilbert-space basis would be admissible but we expect that a Coulomb-Sturmian (CS) basis [13] would be the most advantageous in practical calculations. The reason is that such a basis allows for an exact analytic calculation of the two-body Green's operators and of other two-body quantities [14]. As a result the contour integral can be performed also in the

practice. The practicability of this approach has been demonstrated already in [3–5]. The present paper provides the formalism for extending the previous approach to scattering problems above breakup energies and to repulsive as well as attractive Coulomb-like interactions.

ACKNOWLEDGMENTS

The authors are thankful to Prof. L. D. Faddeev and Prof. O. A. Yakubovsky for fruitful discussions, to Prof. I. V. Komarov for valuable comments and to Prof. W. Plessas for critical reading of the manuscript.

This work has been supported by Hungarian OTKA Grant under Contracts No T026233 and T029003, partially by the Austrian-Hungarian Scientific-Technical Cooperation within project A-14/1998 as well as by Russian Foundation for Basic Research Grant No 98-02-18190 and by RFBR-INTAS Grant No 95-0414.

-
- [1] L. D. Faddeev and S. P. Merkuriev, *Quantum Scattering Theory for Several Particle Systems*, (Kluwer, Dordrech, 1993).
 - [2] J. Noble, J., Phys. Rev. **161**, 945 (1967).
 - [3] Z. Papp and W. Plessas, Phys. Rev. C **54**, 50 (1996).
 - [4] Z. Papp, Phys. Rev. C **55**, 1080 (1997).
 - [5] Z. Papp, Few-Body Systems **24**, 263 (1998).
 - [6] L. D. Faddeev, *Mathematical Aspects of the Three-Body Problem in Quantum Scattering Theory*, Israel Program for Scientific Translation, Jerusalem, (1965).
 - [7] S. P. Merkuriev, Zapiski Nauchnyh Seminarov LOMI, **77**, 148 (1978) (in Russian).
 - [8] S. P. Merkuriev, Ann. Phys. (N.Y.) **130**, 395 (1980); Acta Physica Austriaca, Supplementum XXIII, 65 (1981).
 - [9] V. Vanzani, *Few-Body Nuclear Physics*, (IAEA Vienna), 57 (1978).
 - [10] W. Glöckle, Nucl. Phys. **A141**, 620 (1970); *ibid.* **A158**, 257 (1970).
 - [11] S. L. Yakovlev, Teor. Mat. Fiz. **107**, 531 (1996) (in Russian) [Theor. Math. Phys. **107**, 835 (1996) (in English)].
 - [12] W. Sandhas, *Few-Body Nuclear Physics*, (IAEA Vienna), 3 (1978).
 - [13] M. Rotenberg, Ann. Phys. (N.Y.) **19**, 262 (1962); M. Rotenberg, Adv. Atom. Mol. Phys. **6**, 233 (1970).
 - [14] Z. Papp, J. Phys. A **20**, 153 (1987); Phys. Rev. C **38**, 2457 (1988); Phys. Rev. A **46**, 4437 (1992); Comp. Phys. Comm. **70**, 426 (1992); *ibid.* **70**, 435 (1992); B. Kónya, G. Lévai, and Z. Papp, J. Math. Phys. **38**, 4832 (1997).




FIG. 1. Analytic structure of $g_{x\alpha}(z-z') g_{y\alpha}(z')$ as a function of z' with $z = E + i\epsilon$, $E > 0$, $\epsilon > 0$. The contour C encircles the continuous spectrum of $g_{y\alpha}$ and avoids the singularities of $g_{x\alpha}$. A part of it, which belongs to the second Riemann sheet of $g_{y\alpha}$, is drawn as a broken line.